arXiv:astro-ph/0404202v2 8 May 2004

A new alternative model to dark energy

Yungui Gong
Institute of Applied Physics and College of Electronic Engineering
Chongqing University of Post and Communication
Chongqing 400065, China
gongyg@cqupt.edu.cn

Xi-Ming Chen
College of Electronic Engineering
Chongqing University of Post and Communication
Chongqing 400065, China
chenxm@cqupt.edu.cn

Chang-Kui Duan
Institute of Applied Physics and College of Electronic Engineering
Chongqing University of Post and Communication
Chongqing 400065, China
duanck@cqupt.edu.cn

The recent observations of type Ia supernovae strongly support that the universe is accelerating now and decelerated in the recent past. This may be the evidence of the breakdown of the standard Friedmann equation. Instead of a linear function of the matter density, we consider a general function of the matter density to modify the Freidmann equation. We propose a new model which explains the recent acceleration and the past deceleration. Furthermore, the new model also gives a decelerated universe in the future. The new model gives $\Omega_{\rm m0} = 0.46$ and $z_{\rm T} = 0.44$.

Keywords: dark energy; alternative model; Type Ia supernova.

PACS Nos.: 98.80.-k, 98.80.Es,04.50.+h,14.80.-j

The possible discovery of an accelerating universe from observations of Type Ia supernovae (SNe Ia) leads to a new wave of interest in cosmology [1-4]. Observational results also provide the evidence of a decelerated universe in the recent past [5,6]. On the other hand, the cosmic microwave background (CMB) observations indicate that the universe is spatially flat as predicted by inflationary models [7,8]. These observations suggest that the universe is dominated by dark energy. One simple candidate of dark energy is the cosmological constant. However, the unusual small value of the cosmological constant leads to the search for dynamical dark energy models although cold dark matter cosmological constant (Λ-CDM) models are consistent with the current observations. The quintessence model is one of the alternatives [9-15]. But there is no direct evidence of dark energy and the nature of dark energy remains mysterious. One logical alternative is that the standard Friedmann equation may need to be modified. In this alternative scenario, the universe is just dominated by ordinary pressureless matter, but the laws of gravity and the standard Friedmann equations are modified. The idea of modifying the laws of gravity is not new. The modified Newtonian Dynamics (MOND) was first used to explain the rotation curve in place of the dark matter [16,17]. In MOND, the Newtonian gravity M/r^2 is replaced with $M/r^2+\sqrt{M}/r$. Since M/r^2 gives the standard Freidmann equation $H^2\sim\rho$, $M/r^2+\sqrt{M}/r$ may provide a modified Friedmann equation $H^2\sim\rho^{2/3}\ln\rho+\rho^{2/3}$ [17]. Recall that the brane cosmology gives a non-standard Friedmann equation $H^2\sim\rho+\rho^2$ [18]. Along this line of reasoning, Freese and Lewis recently proposed the Cardassian expansion in which the universe is dominated by the ordinary matter and the Friedmann equation becomes $H^2\sim\rho+\rho^n$ [19,20]. The Cardassian expansion model was later generalized to a more general Friedmann equation $H^2\sim g(\rho)$ [19-25]. In addition, several authors modified the Friedmann equation as $H^2+H^\alpha\sim\rho$ motivated by theories with extra dimensions [21]. Furthermore, Chung and Freese argued that almost any relationship between H and ρ is possible if our universe is a three brane embedded in five dimensional spacetime [19]. In this paper, we discuss an extra-dimension inspired model with generalized Friedmann equation. We first review three models analyzed by Gong and Duan [25]: a model which is equivalent to the generalized Chaplygin gas model [26] in terms of dynamical evolution, the generalized Cardassian model, and the DGP model [21]. Then we propose our new model and explore the property of this new model.

For a spatially flat, isotropic and homogeneous universe with both an ordinary pressureless dust matter and a minimally coupled scalar field Q sources, the Friedmann equations are

$$H^2=\left(\frac{\dot a}{a}\right)^2=\frac{8\pi G}{3}(\rho_{\rm m}+\rho_{\rm Q}), \tag{1}$$

$$\frac{\ddot a}{a}=-\frac{4\pi G}{3}(\rho_{\rm m}+\rho_{\rm Q}+3p_{\rm Q}), \tag{2}$$

$$\dot{\rho_{\rm Q}}+3H(\rho_{\rm Q}+p_{\rm Q})=0, \tag{3}$$

where dot means derivative with respect to time, $\rho_{\rm m}=\rho_{\rm m0}(a_0/a)^3$ is the matter energy density, a subscript 0 means the value of the variable at present time, $\rho_{\rm Q}=\dot Q^2/2+V(Q)$, $p_{\rm Q}=\dot Q^2/2-V(Q)$ and $V(Q)$ is the potential of the quintessence field. The modified Friedmann equations for a spatially flat universe are

$$H^2=H_0^2 g(x), \tag{4}$$

$$\frac{\ddot a}{a}=H_0^2 g(x)-\frac{3H_0^2 x}{2}g'(x)\left(\frac{\rho+p}{\rho}\right), \tag{5}$$

$$\dot\rho+3H(\rho+p)=0, \tag{6}$$

where $x=\Omega_{\rm m}=8\pi G\rho/3H_0^2=x_0(1+z)^3$ during the matter dominated epoch, $1+z=a_0/a$ is the redshift parameter, $g(x)=x+\cdots$ is a general function of x and $g'(x)=dg(x)/dx$. Note that the universe did not start to accelerate when the other nonlinear terms in $g(x)$ started to dominate. For the matter dominated flat universe, $\rho = \rho_{\rm m}$ and $p = p_{\rm m} = 0$, we have $x_0 = \Omega_{\rm m0}$, $g(x_0) = 1$ and $x = \Omega_{\rm m0}(1+z)^3$. To compare the modified model with dark energy model, we make the following identification by using Eqs. (1)-(6)

$$\omega_{\rm Q} = \frac{xg'(x) - g(x)}{g(x) - x}. \tag{7}$$

In general, $\omega_{\rm Q}$ is not a constant. The transition from deceleration to acceleration happens when the deceleration parameter $q = -\ddot{a}/aH^2 = 0$. From Eqs. (4) and (5), we have

$$g[\Omega_{\rm m0}(1+z_{q=0})^3] = \frac{3}{2}\Omega_{\rm m0}(1+z_{q=0})^3 g'[\Omega_{\rm m0}(1+z_{q=0})^3], \tag{8}$$

$$q_0 = \frac{3}{2}\Omega_{\rm m0} g'(\Omega_{\rm m0}) - 1. \tag{9}$$

Gong and Duan analyzed three models by using the WMAP and supernovae data [25]. The first model is

$$g(x) = x + (1 - \Omega_{\rm m0})[A_s + (1 - A_s)(x/\Omega_{\rm m0})^\beta]^{1/\beta},$$

which is equivalent to the generalized Chaplygin gas model $p_c = -A/\rho_c^\alpha$ as dark energy with $\beta = 1 + \alpha$ and $A_s = (8\pi G/3H_0^2(1-\Omega_{\rm m0}))^\beta A$ in terms of the dynamical evolution of the universe. The second model is the generalized Cardassian model

$$g(x) = x[1 + Bx^{\alpha(n-1)}]^{1/\alpha},$$

where $B = (\Omega_{\rm m0}^{-\alpha} - 1)/\Omega_{\rm m0}^{\alpha(n-1)}$, $\alpha > 0$ and $n < 1 - 1/3(1 - \Omega_{\rm m0}^\alpha)$. The third model is the DGP model $g(x) = [a + \sqrt{a^2 + x}]^2$, where $a = (1 - \Omega_{\rm m0})/2$.

Now let us consider a new extra-dimension inspired model $g(x) = x(1 + e^{-\alpha x})^n$ with $\alpha = -\ln(\Omega_{\rm m0}^{-1/n} - 1)/\Omega_{\rm m0} > 0$. At high redshift z, x is very large and $g(x) \sim x$. Therefore the standard model is recovered at early times. During the matter dominated epoch, $p = 0$. Eq. (5) gives

$$\frac{\ddot{a}}{aH_0^2} = -\frac{1}{2}x(1 + e^{-\alpha x})^n + \frac{3}{2}n\alpha x^2(1 + e^{-\alpha x})^{n-1}e^{-\alpha x}. \tag{10}$$

To have a clear picture of this model, we plot the above function for different values of α and n. From figure 1, it is clear that the expansion of the universe slowed down until the recent past, then the expansion speeds up. However, there is an important new feature of this model, the universe will not expand with eternal acceleration. Therefore there is no future event horizon problem. We can also see from figure 1 that the standard expansion recovered at around $z \sim 2$. If we would like to make a comparison with dark energy model, then we get from Eq. (7)

$$\omega_{\rm Q0} = \frac{n(1 - \Omega_{\rm m0}^{1/n})\ln(\Omega_{\rm m0}^{-1/n} - 1)}{\Omega_{\rm m0}(1 - \Omega_{\rm m0})}.$$

In order to see the observational effect of this model, we use the Wilkinson Microwave Anisotropy Probe (WMAP) data [8] and the supernova (SN) data compiled



Fig. 1. $\ddot{a}/aH_0^2$ versus the redshift z for different $\Omega_{\rm m0}$ and n

by Riess *et al.* [4] to fit the model. The parameters $\Omega_{\rm m0}$ and n are determined by minimizing

$$\chi^2 = \sum_i \frac{[\mu_{\rm obs}(z_i) - \mu(z_i)]^2}{\sigma_i^2} + \frac{(\mathcal{R}_{\rm obs} - \mathcal{R})^2}{\sigma_{\mathcal{R}}^2}, \tag{11}$$

where σ_i is the total uncertainty in the SN observation and $\sigma_{\mathcal{R}}$ is the uncertainty in $\mathcal{R}$, the extinction-corrected distance moduli $\mu(z) = 5\log_{10}(d_{\rm L}(z)/{\rm Mpc}) + 25$, $d_{\rm L}(z) = c(1+z)\int_{\rm T}^{t_0} dt'/a(t')$, the CMB shift parameter $\mathcal{R} \equiv \Omega_{\rm m0}^{1/2} H_0 d_{\rm L}(z_{\rm ls})/(c(1+z_{\rm ls})) = 1.716 \pm 0.062$ [27] and $z_{\rm ls} = 1089 \pm 1$ [8]. The best fit parameters to the gold sample SN data in Ref. 4 and the WMAP data are $\Omega_{\rm m0} = 0.46^{+0.09}_{-0.11}$ and $n = 2.6^{+4.7}_{-0.5}$ with $\chi^2 = 181.4$. The $\Omega_{\rm m0}$ and n 1σ confidence contour is shown in figure 2. At 99.5% confidence level, we have $n \geq 1.7$. By using the best fit parameters $\Omega_{\rm m0} = 0.46$ and $n = 2.6$, we find that $\omega_{\rm Q0} = -2.85$ and the transition redshift $z_{\rm T} = 0.44$. From the dynamical Eq. (7) of $\omega_{\rm Q}$, we see that $\omega_{\rm Q} \approx -n\alpha$ when the redshift z is large. The current matter density and the transition redshift $z_{\rm T} \sim 0.5$ are consistent with the constraints obtained by some model independent results [4,14,28].

In conclusion, we propose a new model which explains the past deceleration and the recent acceleration and recovers the standard model at early times. In addition, the new model predicts that the expansion of the universe will slow down in the future.

Acknowledgments

The authors thank the anonymous referee for fruitful comments. The author YG thanks G. Calcagni for pointing out his work. The work is supported by Chongqing University of Post and Telecommunication under grants A2003-54 and A2004-05.



Fig. 2. The left panel shows the 1σ confidence contour for Ω_{m0} and n and the right panel shows the observational $\mu(z)$ plot and the theoretical $\mu(z)$ plot by using the best fit parameters $\Omega_{\mathrm{m0}} = 0.46$ and $n = 2.6$.

References

1. S. Perlmutter *et al.*, Nature **391**, 51 (1998); Astrophys. J. **517**, 565 (1999); P.M. Garnavich *et al.*, Astrophys. J. Lett. **493**, L53 (1998); A.G. Riess *et al.*, Astron. J. **116**, 1009 (1998).
2. J.L. Tonry *et al.*, Astrophys. J. **594**, 1 (2003); B.J. Barris *et al.*, Astrophys. J. **602**, 571 (2004).
3. R.A. Knop *et al.*, astro-ph/0309368.
4. A.G. Riess *et al.*, astro-ph/0402512.
5. A.G. Riess, Astrophys. J. **560**, 49 (2001).
6. M.S. Turner and A.G. Riess, Astrophys. J. **569**, 18 (2002).
7. P. de Bernardis *et al.*, Nature **404**, 955 (2000); Hanany S *et al.*, Astrophys. J. Lett. **545**, L5 (2000).
8. C.L. Bennett *et al.*, Astrophys. J. Supp. Ser. **148**, 1 (2003); D.N. Spergel *et al.*, Astrophys. J. Supp. Ser. **148**, 175 (2003).
9. P.J.E. Peebles and B. Ratra Astrophys. J. Lett. **325**, L17 (1988); B. Ratra and P.J.E. Peebles, Phys. Rev. D **37**, 3406 (1988); C. Wetterich, Nucl. Phys. B **302**, 668 (1988); P.J.E. Peebles and B. Ratra, Rev. Mod. Phys. **75**, 559 (2003).
10. R.R. Caldwell, R. Dave and P.J. Steinhardt, Phys. Rev. Lett. **80**, 1582 (1998); I. Zlatev, L. Wang and P.J. Steinhardt, Phys. Rev. Lett. **82**, 896 (1999).
11. P.G. Ferreira and M. Joyce, Phys. Rev. Lett. **79**, 4740 (1997); Phys. Rev. D **58**, 023503 (1998); S. Perlmutter, M.S. Turner and M. White, Phys. Rev. Lett. **83**, 670 (1999).
12. V. Sahni and A.A. Starobinsky, Int. J. Mod. Phys. D **9**, 373 (2000); C. Rubano and J.D. Barrow, Phys. Rev. D **64**, 127301 (2001).
13. A.A. Sen and S. Sethi, Phys. Lett. B **532**, 159 (2002); Y. Gong, Class. Quantum Grav. **19** 4537 (2002).
14. U. Alam, V. Sahni, T.D. Saini and A.A. Starobinsky, astro-ph/0311364; U. Alam, V. Sahni and A.A. Starobinsky, astro-ph/0403687.
15. T. Padmanabhan and T.R. Choudhury, Phys. Rev. D **66**, 081301 (2002) ; J.S.

Bagla, H.K. Jassal and T. Padmanabhan, Phys. Rev. D **67**, 063504 (2003); T. Padmanabhan, Phys. Rep. **380**, 235 (2003); T. Padmanabhan and T.R. Choudhury, Mon. Not. Roy. Astron. Soc. **344**, 823 (2003).
16. M. Milgrom, Astrophys. J. **270**, 365 (1983).
17. A. Lue and G.D. Starkman, astro-ph/0310005.
18. P. Binétruy, C. Deffayet and D. Langlois, Nucl. Phys. B **565**, 269 (2000); Y. Gong, gr-qc/0005075.
19. D.J. Chung and K. Freese, Phys. Rev. D **61**, 023511 (1999).
20. K. Freese and M. Lewis, Phys. Lett. B **540**, 1 (2002); K. Freese, Nucl. Phys. Suppl. Ser. **124**, 50 (2003); P. Gondolo and K. Freese, Phys. Rev. D **68**, 063509 (2003).
21. G.R. Dvali, G. Gabadadze and M. Porrati, Phys. Lett. B **485**, 208 (2000); C. Deffayet, Phys. Lett. B **502**, 199 (2001); C. Deffayet, G.R. Dvali and G. Gabadadze, Phys. Rev. D **65**, 044023 (2002); G.R. Dvali and M. Turner, astro-ph/0301510; A. Lue, R. Scoccimarro and G. Starkman , Phys. Rev. D **69**, 044005 (2004).
22. S. Sen and A.A. Sen, Astrophys. J. **588**, 1 (2003) ; A.A. Sen and S. Sen, Phys. Rev. D **68**, 023513 (2003); M. Szydlowski and W. Czaja, astro-ph/0309191; W. Godlowski, M. Szydlowski and A. Karweic, astro-ph/0309569; G. Calcagni, hep-ph/0402126.
23. Z.H. Zhu and M. Fujimoto, Astrophys. J. **581**, 1 (2003); **585**, 52 (2003); Astrophys. J. **602**, 12 (2004); Z.H. Zhu , M. Fujimoto and X.T. He, Astrophys. J. **603**, 365 (2004).
24. Y. Wang, K. Freese, P. Gondolo and M. Lewis, Astrophys. J. **594**, 25 (2003); T. Multamaki, E. Gaztanaga and M. Manera, Mon. Not. Roy. Astron. Soc. **344**, 761 (2003); A. Dev, J.S. Alcaniz and D. Jain, astro-ph/0305068; W.J. Frith, Mon. Not. Roy. Astron. Soc. **348**, 916 (2004); S. Nesseris and L. Perivolaropoulos, astro-ph/0401556.
25. Y. Gong and C.K. Duan, gr-qc/0311060; astro-ph/0401530; Y. Gong and X.M. Chen, gr-qc/0402031.
26. A. Kamenshchik, U. Moschella and V. Pasquier, Phys. Lett. B **511**, 265 (2001); N. Bilic, G.G. Tupper and R.D. Viollier, Phys. Lett. B **535**, 17 (2002); M.C. Bento, O. Bertolami and A.A. Sen, Phys. Rev. D **66**, 043507 (2002); D. Carturan and F. Finelli, Phys. Rev. D **68**, 103501 (2003); L. Amendola, F. Finelli, C. Burigana and D. Carturan, J. Cosmology Astroparticle Phys. **0307**, 005 (2003); J.V. Cunha, J.S. Alcaniz and J.A.S. Lima, Phys. Rev. D **69**, 083501 (2004).
27. J.R. Bond, G. Efstathiou and M. Tegmark, Mon. Not. Roy. Astron. Soc. **291**, L33 (1997); A. Melchiorri, L. Mersini, C.J. Ödman and M. Trodden, Phys. Rev. D **68**, 043509 (2003); Y. Wang and P. Mukherjee, astro-ph/0312192; Y. Wang and M. Tegmark, astro-ph/0403292.
28. R.A. Daly and S.G. Djorgovski, Astrophys. J. **597**, 9 (2003); astro-ph/0403664; Y. Gong, astro-ph/0401207.